

27 figures

25/1




FIGURE 1

FIGURE 2A

FIG. 2B




FIGURE 2B




FIGURE 3

FIG. 4




FIGURE 4

FIGURE 5




FIGURE 5





FIGURE 6



1. *Staphylococcus aureus* (ATCC 12228) was grown in tryptic soy broth (TSB) (Difco) supplemented with 0.5% yeast extract (Difco) and 0.5% glucose (Difco) at 37°C. Cells were harvested at mid-log phase (OD₆₀₀ = 0.5) and washed with phosphate buffered saline (PBS) (pH 7.4) containing 100 μg/ml penicillin, 100 μg/ml streptomycin, and 100 μg/ml nystatin. Cells were then washed with PBS containing 100 μg/ml penicillin, 100 μg/ml streptomycin, and 100 μg/ml nystatin. Cells were then washed with PBS containing 100 μg/ml penicillin, 100 μg/ml streptomycin, and 100 μg/ml nystatin.

FIG. 9




FIGURE 9




FIGURE 8




FIGURE 10




FIGURE 11




FIGURE 12

202500054600




FIGURE 13




FIGURE 14




FIGURE 15




FIGURE 16

1. 1992-1993
 2. 1994-1995
 3. 1996-1997




FIGURE 17




FIGURE 18




FIGURE 19A




FIGURE 19B




FIGURE 19C




FIGURE 19D




FIGURE 19D





FIGURE 19E



■■■■■ ■ PUMP PRESSURE
 ■■■■■ ■ RETURN PRESSURE

FIGURE 19F